



A service of the National Library of Medicine  
and the National Institutes of Health

My NCBI  
[Sign In] [Reg]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Bc

Search PubMed for pulse cell cycle metaphase [Preview] [Go] [C]

Limits Preview/Index History Clipboard Details

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorials

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI

Related Resources

Order Documents

NLM Mobile

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

- Search History will be lost after eight hours of inactivity.
- Search numbers may not be continuous; all searches are represented.
- To save search indefinitely, click query # and select Save in My NCBI.
- To combine searches use #search, e.g., #2 AND #3 or click query # for more options.

Search	Most Recent Queries	Time	Result
#19	Search pulse cell cycle metaphase	10:53:22	76
#14	Search electrical cell cycle metaphase	10:51:20	22
#13	Search electrical cell cycle	10:46:18	480
#12	Search synchronize voltage	10:45:55	38
#11	Search synchronize metaphase voltage	10:45:37	0
#8	Search gsk3 apoptosis	05:58:09	44
#7	Search gsk3 angiogenesis	05:56:57	0
#4	Search gsk3b	05:56:47	8
#3	Search gsk3	05:54:56	388
#1	Search gsk3 endothelial	05:54:13	4

Clear History

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Sep 18 2006 12:27:35

cell became confluent, it was treated with a 0.05-0.53 M trypsin EDTA solution for 5 minutes to extricate it from the cell, and it was diluted with 0.4 ml HBS buffer solution and electrical stimulation was applied.

5           The electrical stimulation was applied under conditions of a voltage of 266 V, a condenser value of 975  $\mu$ F, and a resistance value of infinity, for 20 msec. The embryonic stem cell was placed on ice for 10 minutes immediately after electrical stimulation, and then seeded on a feeder cell in a dish with a diameter of 60 mm to  
10   culture.

1-2. Synchronous culture of embryonic stem cell to metaphase stage of cell division

The embryo stem cells were initially cultured with feeder cells after applying electrical stimulation, and 3 days after initial  
15   culture they were cultured in a culture liquid to which nocodazole (0.4  $\mu$ g/ml) was added, for 2-4 hours. Cells in the metaphase stage lose their adhesive property and rise up in the culture liquid. Therefore, the upper part of the culture liquid was recovered and centrifuged at 1000 rpm for 5 minutes to obtain embryonic stem cells of the  
20   metaphase stage.

1-3. Nuclear transfer and artificial culture

To a female B6CBF1(C57BL/6 x CBA) mouse, PMSG (pregnant mare serum gonadotropin) and hCG (human chorionic gonadotropin) were respectively administrated with an interval of 48

(Table 1)

	No. of nuclear transferred embryos cultured	No. of blastocysts transferred	No. pregnant / no. of recipients (%)	No. of implantation sites-(%)	No. of live pups(%)
Comparative Example 1	320	148(46.3)	5/12(41.7)	24(16.2)	0(0)
Example 1	245	110(44.9)	6/9(66.7)	57(51.9)	3(2.7)

As can be seen from the Table 1, Example 1 wherein electrical stimulation was applied shows higher implantation and productivity rates than Comparative Example 1.

(2) Examination of productivity according to differentiation of embryonic stem cell

In order to examine productivity according to differentiation of an embryonic stem cell used as a donor cell, no. of implantation sites, implantation rate, and no. of live pups of the reconstituted eggs according to Example 3 and Comparative Example 2 were measured and results shown in the following Table 2.

(Table 2)

	No. of nuclear transferred embryos cultured	No. of blastocysts transferred (%)	No. of implantation sites (%)	No. of live pups (%)
Comparative Example 2	55	37(67)	13(24)	0(0)
Example 3	70	51(73)	39(56)	3(5.9)

As seen from Table 2, Example 3 using an undifferentiated embryonic stem cell as a donor cell shows a high blastocyst stage